

Mass developments of a small sized ecotype of *Arthrospira fusiformis* in Lake Ololdien, Kenya, a new feeding ground for Lesser Flamingos in East Africa

Lothar KRIENITZ^{1*}, Pawan K. DADHEECH^{1,2,3} & Kiplagat KOTUT⁴

¹Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Alte Fischerhütte 2, D-16775 Stechlin, Germany; * Corresponding author e-mail: krie@igb-berlin.de

²Central University of Rajasthan, Department of Microbiology, Bandarsindri 305801, Kishangargh, India

³Government College, Department of Botany, Ajmer 305001, Rajasthan, India

⁴Embu University College, P.O. Box 6-60100 Embu, Kenya

Abstract: The Lesser Flamingos constantly commute between the soda lakes of East Africa in search of food. Their preferred food, the cyanobacterium *Arthrospira fusiformis*, usually establishes dense populations in saline-alkaline habitats. However, the algal food resources are not stable and crash unexpectedly from time to time. Hence any site that provides food in suitable quality and quantity makes a valuable contribution to the survival of these nearly endangered birds. This paper reports on the development of a new feeding ground for the Lesser Flamingos; Lake Ololdien, a former bay of the freshwater lake Naivasha. Phytoplankton succession and biomass, salinity and flamingo numbers were analysed over the period 2001–2013. As a result of a progressive increase in salinity to values of between 3 and 6 ppt, Ololdien has since 2006 hosted dense blooms of a small sized ecotype of *A. fusiformis*. Molecular phylogenetic analyses using the 16S–23S internal transcribed spacer (ITS) and partial sequences of the beta and alpha subunits including intergenic spacer (*cpcBA*–IGS) of the phycocyanin operon confirmed that a field clone and one isolated strain of *Arthrospira* from Lake Ololdien was similar to the strains in other soda lakes of Kenya. PCR-based results confirmed the non-toxic character of the isolated *Arthrospira* strain. The biomass of *Arthrospira* in the whole lake ranged from 236 to 301 t dry weight in the phase when more than 100,000 flamingos were present (2006–2012). Based on the biomass of *Arthrospira* present, the carrying capacity of Lake Ololdien was calculated to be 150,000–250,000 Lesser Flamingos. In the phase of low flamingo numbers (2013), the biomass of *Arthrospira* in the lake rose to a value of 922 t.

Key words: *Arthrospira*; cyanobacteria; Lake Naivasha; Lake Ololdien; Lesser Flamingo; salinity

INTRODUCTION

The African saline alkaline lakes are hosts to two species of flamingos, the Greater Flamingo (*Phoenicopterus roseus* PALLAS) and the Lesser Flamingo (*Phoeniconaias minor* SAINT HILAIRE). The two flamingo species differ in their food preference. Whereas the Greater Flamingo is a generalist that consumes small planktonic and benthic animals such as molluscs and arthropods, as well as mud and algae, the Lesser Flamingo is highly specialised as it depends on cyanobacteria and algae (JENKIN 1957; BROWN 1959). Its preferred food is the cyanobacterium *Arthrospira fusiformis* (WORONICHIN) KOMÁREK et J.W.G. LUND (often erroneously named as “*Spirulina*”) (VARESCHI 1978). However, this algal food resource is not stable and crashes periodically (VARESCHI & JACOBS 1985). Alternatively, Lesser Flamingos can feed on benthic diatoms and other eukaryotic algae. However, the grow rate and available biomass of these alternative food sources are much lower (TUIE 1981, 2000). Lesser Flamingos are mainly distributed in the soda lakes of East Africa, where about 2 million individuals

of this enigmatic bird have been counted (CHILDRESS et al. 2008). Rising demographic pressure in the lakes’ catchments has led to an accelerated degradation of the soda lakes. The consequence of these changes has been the alteration of the feeding grounds of Lesser Flamingo leading to a decline in the amount of food cyanobacteria of high nutritional value (KRIENITZ & KOTUT 2010). The special diet requirement and the need for undisturbed breeding habitats have made the Lesser Flamingo very sensitive to environmental changes. The only site in East Africa where this bird can breed successfully is Lake Natron in Tanzania. In its search for adequate food and suitable breeding places, the Lesser Flamingos exhibit an unsettled behaviour. Satellite tracking of flamingos fitted with transmitters in 2001 and 2002 showed that the longest distance made by one of the flamingos in a period of 15 months was 7,870 km. In this period the flamingo made a total of 70 stopovers at 11 different lakes of East Africa (CHILDRESS et al. 2006).

The major threats to the survival of Lesser Flamingos are the loss or deterioration of their unique habitats. In the red list of the International Union for the Conservation of Nature (IUCN), this species is

classified as nearly threatened. An International Single Species Action Plan for Conservation of the Lesser Flamingo was established to protect this character bird of harsh soda lakes (CHILDRESS et al. 2008).

In this paper, we report on the development of a new feeding ground for Lesser Flamingos; Lake Oloidien, a former bay of Lake Naivasha in Kenya. The negative effects of overexploitation and degradation of the Lake Naivasha freshwater ecosystem (HARPER et al. 2002, 2011), has been accompanied by a positive development: a flamingo–lake was born (Fig. 1). The ambivalent development characterized by increasing salinity and a switch in the phytoplankton composition created favourable conditions for the flamingos (HARPER et al. 2006; KRIENITZ & KOTUT 2010; KRIENITZ et al. 2013). The focus of this study was the description and characterization of the food source and its suitability for Lesser Flamingos in Lake Oloidien. In this paper, the carrying capacity of the lake was computed and the findings used to evaluate whether the estimated high numbers of flamingos could overexploit the *Arthrospira* food resource in the lake. Molecular analyses were carried out to characterize the phylogenetic position of the *Arthrospira* population from Oloidien and its possible toxicity in comparison to other flamingo lakes in the region.

MATERIAL AND METHODS

Study site, sampling and microscopic studies of the phytoplankton. Lake Oloidien is located in the Great African Rift Valley next to Lake Naivasha, about 80 km North of Nairobi at an altitude of 1890 m above sea level (Fig. 2). Paleolimnological studies covering a time span of ~ 1870 – 1991 revealed that the water level of the lake fluctuated between 19 and 4 m while conductivity varied between ~ 250 and 14,000 $\mu\text{S}\cdot\text{cm}^{-1}$ (VERSCHUREN et al. 2000). Consequently, lakes Naivasha and Oloidien experienced periods of being one or separate water bodies. The present situation dates back to the 1980s when decreasing lake levels led to a separation of Lake Oloidien from Lake Naivasha. The conductivity at this time was 660 μS (KALFF & WATSON 1986). In the period 2001–2005, a conductivity of 3890 to 5270 $\mu\text{S}\cdot\text{cm}^{-1}$ was measured (BALLOT et al. 2009). The lake has no surface inflow and no surface outlet. At present, the endorheic basin has a surface area of about 450 ha and a maximum depth of 7 m. The basin of Lake Oloidien is a caldera of a crater with the shape of a truncated cone. Based on its morphometric features, we calculated its volume to be $1.26 \times 10^7 \text{ m}^3$. The annual rainfall in this area is 500 – 700 mm and is the main source of water. Lake Oloidien is highly eutrophic with total phosphorus and total nitrogen values of between 0.4 and 1.0 $\text{mg}\cdot\text{l}^{-1}$, and 0.9 and 6.3 $\text{mg}\cdot\text{l}^{-1}$ respectively (BALLOT et al. 2009; our own measurements). The pH values increased from 9.3 at the beginning of the study to 10.2 – 10.4 in the last five years.

The main sampling point (1) at Oloidien (Fig. 3) is located at the western shore near the village of Kongoni



Fig. 1. Lesser Flamingos at Lake Oloidien. Although the birds were building nests, breeding was not possible because of disturbance by humans and predators.

(00°49'00"S, 36°15'50"E). During the period 2002–2013, 21 sampling trips were carried out at irregular intervals. The samples were taken from a depth of about 10 cm below the water surface and fixed with Lugol's solution for phytoplankton counting. Water samples for morphological studies were concentrated with a plankton net of 25 µm mesh size and fixed with formaldehyde. During the sampling expeditions conducted in April 2012, November 2012, and January 2013, phytoplankton samples were also collected from eight different sampling points in addition to the sample from the main sampling point (Fig. 3). Samples were drawn from different depths (0, 2, 4, and 6 m) using a Ruttner sampler. All the samples were counted in sedimentation chambers (Hydro-Bios Apparatebau GmbH, Kiel, Germany) under an inverted microscope Eclipse TS 100 (Nikon Corporation, Tokyo, Japan) following the method of UTERMÖHL (1958). The phytoplankton biomass (fresh weight, FW) was calculated by geometric approximations using the computerized counting programme OPTICOUNT (OPTICOUNT 2008). The specific density of phytoplankton cells was taken as 1 g.cm⁻³. Phytoplankton was photographically documented under a Nikon Eclipse E 600 light microscope using a Nikon digital camera DS-Fi1, and Nikon software NIS-Elements D (Nikon Corporation, Tokyo, Japan). The size and morphology of *Arthrospira* from Oloidien was compared with those from lakes Nakuru and Bogoria. In addition to FW measurement in Lake Oloidien, we also computed dry weights (DW) of *Arthrospira* using samples collected in the last ten sampling trips and carried out in the period 2006–2013 when *Arthrospira* was the dominant phytoplankton. For this purpose, 100 ml of a fresh water sample was filtered through pre-weighed glass fibre filters (Whatman GF/C, Whatman International Ltd Maidstone, England) in triplicate after which the filters were dried. After drying, the filters with samples were weighed again and the DW of the phytoplankton in 100 ml of sample computed. A polycarbonate filter holder SM 16510 (Sartorius AG, Göttingen, Germany) and a manual vacuum pump VacuMan (Bürkle GmbH, Bad Bellingen, Germany) were used as filtration apparatus in the field. For the molecular analyses of field samples, 1 l of the fresh sample was filtered using membrane filters with a pore-size of 0.6 µm (Schleicher & Schuell GmbH, Dassel, Germany). The salinity was measured at each sampling date using a WTW Multiline P4 meter (Wissenschaftlich Technische Werkstätten Weilheim, Germany).

Estimation of flamingo numbers. A visual estimate of the number of flamingos was made and sorted into four categories (> 100,000, >20,000, 20,000–10,000, and <1,000 individuals). These rough estimates were carried out at strategic points at the lake shore and from the boat. In many cases, it was possible to adjust our estimates by comparing our observed findings with those of the boat guides (Oloidien Community Boats, members of the initiative “Friends of Flamingos”).

Cultivation of *Arthrospira*. To establish clonal cultures of *Arthrospira*, single filaments from fresh field samples were isolated with a micro capillary tube and transferred into liquid modified BOURRELLY medium (KRIENITZ & WIRTH 2006). The BOURRELLY medium without salt (B) was mixed with BOURRELLY medium with salt (BS) in distinct proportions to receive a range of salinity (0, 2, 4, 8, 12, and 16 ppt). The BS medium contained 0.3 g Na₂CO₃ and 15 g




Fig. 2. Map showing the Rift Valles lakes in Kenya, modified from Brown (1959).



Fig. 3. Map taken from Google Earth showing Lake Oloidien and the sampling sites.

NaCl per litre. The cyanobacterial suspensions were grown in 100 ml Erlenmeyer flasks at room temperature (23 ± 2 °C) under a 14/10 light/dark regime. The strains isolated by LK were deposited in the algal strain collection of the Leibniz-Institute of Freshwater Ecology and Inland Fisheries at Stechlin. The strain KR2012/1 was used to test the growth at different salinities and molecular studies.

Molecular studies. The collected field samples were scrubbed from dried filter with a sterile scalpel. The genomic DNA of the field sample and the isolated strain KR2012/1 of *Arthrosira* was extracted using Dynabeads DNA DIRECT




Fig. 4. Phytoplankton biomass (FW), succession of main phytoplankton groups, number of flamingos, percentage contribution of *Arthrospira* to total phytoplankton biomass, and salinity in Lake Oloidien during the period 2001–2013.

System I (Invitrogen/Dyanl Biotech, Oslo, Norway) following the steps outlined in the manufacturer's manual. The two genetic loci (ITS and *cpcBA*–IGS) were employed for both cases as these markers are considered the most suitable to characterize *Arthrospira* strains (BAURAIN et al. 2002; DADHEECH et al. 2010). The polymerase chain reaction (PCR) of ITS region and *cpcBA*–IGS locus was performed in a Peltier Thermal Cycler PTC 200 (MJ Research Inc., San Francisco, USA) using PCR cocktail and protocol as previously described (DADHEECH et al. 2010). The primers 16S3'F and B235'R (BAURAIN et al. 2002) were used to amplify ITS region while the primers *cpc_ar*F and *cpc_ar*R (BALLOT et al. 2004) were used to obtain partial sequence of *cpcBA*–IGS. PCR products of the ITS region and *cpcBA*–IGS locus were purified through Qiaquick PCR purification columns (Qiagen, Hilden, Germany). The purified products of field sample were cloned using the Zero Blunt® Topo® PCR cloning kit (Invitrogen, Germany) according to manufacturer's instructions. Positive colonies were selected and amplified with PCR and then sequenced to retrieve the sequence of ITS region and *cpcBA*–IGS locus. The clones of ITS and *cpcBA*–IGS were designated as Olo_arthro_its and Olo_arthro_cpc respectively. Both strands of purified DNA (ITS and *cpcBA*–IGS) were sequenced with the same primers used for the PCR on ABI 3100 Avant Genetic Analyzer (Applied Biosystem, Applera, Darmstadt, Germany).

The genetic potential of the investigated *Arthrospira* sp. strain (KR2012/1) to produce a variety of cyanotoxins was assessed using different primer sets. The primers and PCR protocol for each genetic locus, HEPF/HEPR for microcystin/nodularin (JUNGBLUT & NEILAN 2006), FAA/

RAA for *mcyB* (NEILAN et al. 1999), AnaC–genF/AnaC–genR for anacystin (RANTALA–YLINEN et al. 2011) and sxtA–F/sxtA–R for saxitoxin (AL–TEBRINEH et al. 2010) were employed for amplification.

The sequences of ITS and *cpcBA*–IGS locus of *Arthrospira* particularly from Asian, African and Mexican origin were retrieved from nucleotide NCBI database and aligned with sequences obtained in present study using software MUSCLE (Edgar 2004). Alignment was checked visually using the Manual Sequence Alignment Editor, Align v05/2008 (HEPPERLE 2008). The phylogenetic trees were constructed by the maximum likelihood (ML) method using the program MEGA v5.0 (TAMURA et al. 2011) with default settings, Jukes–Cantor model of nucleotide. Confidence values for the edges of the maximum–likelihood tree were computed by bootstrapping of 1000 replications. The nucleotide sequences reported in this study have been deposited in the NCBI database under the GenBank accession number JX853612–JX853613 (ITS) and JX853614–JX853615 (*cpcBA*–IGS).

RESULTS

In the years 2001–2005, the number of flamingos at Oloidien was very low and never exceeded 1,000 individuals. In 2006, the situation changed dramatically. The number of flamingos increased from less than 1,000 in February to more than 20,000 in September




Fig. 5. Microphotographs of *Arthrospira fusiformis*; for direct comparison, all figures are provided with the same magnification. (a–d) field samples: (a, b) small sized ecotype from Lake Oloidien, (c) Lake Nakuru, (d) Lake Bogoria; (e, f) strain KR 2012/1 isolated from Lake Oloidien as small ecotype which grows in suspension cultures up to larger sized variants. Scale bar 30 μm .

2006 (Fig. 4). In the following years, the flamingo numbers fluctuated between 10,000 and 20,000. Towards the end of 2010, the flamingos established a stable population of up to 50,000. In 2012, there was again an increase in the number of birds with more than 100,000 flamingos being observed. By April 2012, between 150,000 and 250,000 flamingos were recorded in Oloidien. In November 2012 and January 2013, the number of flamingos decreased dramatically to only about 1000.

The rise in flamingo numbers at Oloidien resulted from a shift in the phytoplankton community structure brought about by the considerable changes in water chemistry. Between 2001 and 2005, the water was moderately saline with a salinity of about 2 ppt. During this low salinity phase, the phytoplankton was dominated by coccoid green algae as well as coccoid and colonial cyanobacteria. The filamentous cyanobacterium *Anabaenopsis* was subdominant and reached its maximum in 2006 with densities of 50 – 60 mg.l^{-1} . However, in 2006 the salinity level reached a value of more than 3 ppt. The attainment of this salinity level led to a massive growth of *Arthrospira fusiformis*, the favourite food for the Lesser Flamingo. Because of the long dry seasons in subsequent years,

salinity increased reaching a maximum value of 5.5 ppt in January 2010. Although the heavy rains received in 2010/2011 diluted the water to salinity values of 3 ppt, *Arthrospira* development remained on the same high biomass level of more than 150 mg.l^{-1} FW (Fig. 4). Even in November 2012 and January 2013, when heavy rains resulted in a drop in salinity to values of 2.3 and 2.2 ppm respectively, phytoplankton dominance by *Arthrospira* remained at 100%. Because of the extremely low numbers of flamingos, it was interesting to measure the phytoplankton biomass in this phase when the grazing pressure was insignificant. The *Arthrospira* biomass in Lake Oloidien rose to a maximum of 623 mg.l^{-1} and was entirely made of small filament forms (Fig. 5a). The majority of filaments were 4–5 μm wide with a few being 6–8 μm wide. Filament lengths ranged from 60–100 μm with 2–4 coils. The few broader filaments had lengths of up to 200 μm with 4–6 coils (Fig. 5b). We found only a few gas vesicles in the cells of the small filaments. Spiral diameter in each filament ranged from 25–40 μm . Comparing the morphology of *Arthrospira* from Oloidien with that from the main flamingo lakes of the Kenyan Rift Valley, the form found in Oloidien was considerably smaller than the specimens from Lake Nakuru (Fig. 5c)

(filaments 10–13 μm wide and 150–300 μm long) and Lake Bogoria (Fig. 5d) (filaments 12–15 μm wide and up to 700 μm long, with up to 20 coils). Many gas vesicles are visible in these large filaments. When grown in culture, the strains from Lake Oloidien, lost their small size and reached larger dimensions comparable to those measured in the field material of Nakuru and Bogoria (Fig. 5e, f) within 30 days of growth, even under a salinity of 2 ppt. Laboratory experiments on the biomass production of strain KR2012/1 under different salinity regimes resulted in a maximum yield at 8 ppt salinity. At a salinity of 16 ppt, a considerably lower biomass production was observed while at a salinity of 0.1 ppt, very limited biomass production was recorded (Fig. 6).

The buoyancy behaviour of the *Arthrospira* forms from the three different lakes contrasted sharply. Depth samples drawn from different depths of the water column using a Ruttner sampler revealed that the small form in Oloidien was evenly distributed down the water column and did not show a dense accumulation on the upper layer of the water. In Lake Nakuru, we observed a moderate layering in the upper five centimetres while in Lake Bogoria, a strong accumulation of *Arthrospira* on the water surface was observed leading to dense paste-like scums of the cyanobacterium in the upper 1–2 cm. This phenomenon could be attributed to the comparatively higher number of gas vesicles in the large filaments.

We obtained sequences of 411 bp long for ITS region and of 489 bp long for *cpcBA*–IGS locus. The phylogenetic position of field clone and isolate (KR2012/1) were determined in ML trees of ITS and *cpcBA*–IGS regions. For both loci, *Arthrospira* sequences of the same strain from different Kenyan, Indian and Mexican lakes were also compared to our sequences. The ITS and *cpcBA*–IGS sequences of the field clone as well as the isolate occurred in the cluster comprising strains of *Arthrospira* from several Kenyan lakes (Fig. 7). PCR studies aimed at determining the cyanotoxin production potential by the *Arthrospira* isolate under investigation using several specific primer pairs did not lead to the amplification of any genetic locus responsible for the production of cyanotoxins (data not shown). Tests for the presence of microcystins and anatoxin–a confirmed the absence of toxins in a field sample collected on 21 November 2011 (Jutta Fastner, unpubl. data).

Biomass measurements established that 1 mg DW was equivalent to 8.5 mg FW as reported by VARESCHI (1982). The mean *Arthrospira* biomass obtained in the eight sampling trips carried out in the phase of high abundance of flamingos (September 2006–April 2012) was $153 \pm 21 \text{ mg.l}^{-1} \text{ FW}$ and $18.7 \pm 2.5 \text{ mg.l}^{-1} \text{ DW}$ (standard deviation given) at the main sampling point. *Arthrospira* biomass measurements carried out in nine sampling points on 14 April 2012 and 24 April 2012 gave mean biomass values of $190 \pm$

$15 \text{ mg.l}^{-1} \text{ FW}$ and $22.4 \pm 1.7 \text{ mg.l}^{-1} \text{ DW}$ (14 April 2012) and $203 \pm 13 \text{ mg.l}^{-1} \text{ FW}$ and $23.9 \pm 1.5 \text{ mg.l}^{-1} \text{ DW}$ (24 April 2012). As the *Arthrospira* biomass was found to be evenly distributed down the water column, the total DW for the whole lake was calculated to be 236 t based on mean value for the period 2006–2012 and 301 t for 24 April 2012. In November 2012 and January 2013 when the number of flamingos dropped to about 1000 birds, the grazing pressure on the *Arthrospira* population became insignificant. During this period, the total biomass calculated for the whole lake rose a high of 880 t (November 2012) and 922 t (January 2013).

DISCUSSION

Increase of flamingo numbers at Lake Oloidien

The soda lakes in Kenya host the densest population of flamingos in East Africa. Although the number of birds in each lake varies widely, the total number sometimes rises to more than 2 million individuals (BROWN 1959; CHILDRESS et al. 2008). Under favourable conditions, the main lakes of Nakuru, Bogoria and Elmenteita can accommodate about 1.5 million (BROWN 1959, VARESCHI 1978), 2 million (BROWN 1959; HARPER et al. 2003), and 108,081 (OWINO et al. 2001) birds respectively. Twice a year, estimates of bird numbers in the lakes is carried out by ornithologists through an initiative of the African Waterfowl Census (MAINA & KYUNGU 2009). By compiling the bird numbers simultaneously at all the main distribution areas, an overview of the total population density of a bird species and its trends can be established (NASIRWA et al. 2007). However, because of the itinerant behaviour of Lesser Flamingos, such a counting programme may not detect the peak numbers of flamingos at a given lake. Short term changes in population density are extremely difficult to determine and depend on randomly scheduled counting




Fig. 6. Dry weight of *Arthrospira fusiformis*, strain KR 2012/1, under different salinity.




Fig. 7. (a) Maximum-likelihood phylogenetic tree based on ITS sequences of *Arthrospira*. *Phormidium autumnale* UTCC 579 was used as an outgroup. The tree was derived using Jukes–Cantor model. The numbers above branches indicate bootstrap support (>50%) from ML (1,000) replicates. (b) Maximum-likelihood phylogenetic tree based on *cpcBA*–IGS sequences of *Arthrospira*. *Spirulina subsalsa* PD2002/gea was used as the outgroup. The tree was derived using Jukes–Cantor model. The numbers above branches indicate bootstrap support (>50%) from ML (1,000) replicates.

programmes. At the beginning of the massive flamingo establishment at Lake Oloidien in 2006, Earthwatch research teams gave an estimate of nearly a quarter of a million flamingos in October (Earthwatch Institute 2006). However, a month earlier, we had counted only about 20,000 flamingos. Members of the “Friends of Flamingos” group at Oloidien who undertake daily boat trips on the lake confirmed that during the phase of a rapid increase in flamingo numbers that begun in 2010, the counts have varied widely with daily fluctuations sometimes being by more than 50%. The birders have reported that in January – March 2012, nearly one half of a million Lesser Flamingos collected their food from Lake Oloidien. Enthusiastic calculations posted

in the internet for the same period have given values of about 1.5 millions birds at the lake (STANDARD MEDIA 2012). Our irregular observations over a period of 12 years have clearly demonstrated a progressive increase in numbers of Lesser Flamingos at Lake Oloidien with numbers varying between 100,000 and 250,000 individuals. Owing to an expected strong grazing pressure on the primary producers in the lake by the high numbers of flamingos, we characterized the quality and quantity of the food algae available. At the end of our observation period (November, 2012 and January 2013), the number of flamingos dropped drastically to only about 1000 individuals. It is suspected that they could have moved to Lake Natron for breeding (BBC

NATURE 2012). We used this occasion when the grazing pressure was absent to quantify the food concentration.

Characterization of the small type of *Arthrospira fusiformis* from Lake Oloidien

Arthrospira fusiformis observed in Lake Oloidien belongs to the variety *minor*, which refers to populations of fine filaments 4–5 µm in width (RICH 1932) and whose earlier designation was *A. platensis* (NORDSTEDT) GOMONT, a synonym of *A. fusiformis*. RICH first reported the presence of var. *minor* in the Crater Lake. The Crater Lake Sonachi is located at a distance of only 3.7 km from Lake Oloidien. Interestingly, at the time of RICH's collection in 1929, water salinity in the Crater Lake was about 3 ppt (computed from conductivity data published by VERSCHUREN et al. 1999). Presently, the salinity of this lake is about 10 ppt and its phytoplankton composition is dominated by the large size form of *Arthrospira fusiformis* (BALLOT et al. 2005). Generally, under field conditions, the small variety is rare. With the exception of its dominance in Lake Oloidien, the variety *minor* is rarely observed. We found it at a comparably high density at Kamfers Dam (South Africa), which has a similar salinity range of about 3 ppt (Lothar Krienitz, unpubl. results). Moreover, the small size variety of *A. fusiformis* has been reported to be a contaminant of outdoor mass cultivations of the “normal” form of *Arthrospira* (VONSHAK & RICHMOND 1988). Although the small sized variant of *A. fusiformis* has been found to be more sensitive to high ultraviolet radiation, this deficiency is usually compensated for by the lower cell buoyancy that results in a more-or-less uniform distribution within the water column. This reduces the risk associated with strong solar radiation on the water surface (GAO & MA 2008). The uniform distribution of the small size *Arthrospira* in the water column was a consistent feature of Oloidien throughout the present study. In contrast, the large size form established dense surface scums in Bogoria. In Lake Nakuru, VARESCHI (1982) identified two different types of *Arthrospira* at different periods; a small size type (resembling the var. *minor*) that was uniformly distributed throughout the water column during phases of dense blooms and a large size type that establishes surface scums on the lake during phases of low phytoplankton density. He recognized two different taxa, spec. 1 and spec. 2. However, we were not able to corroborate this differentiation. Our growth experiments under different medium salt concentration revealed that the small size variant of *Arthrospira* does well under low to moderate salinity conditions. Data from past studies however suggest a high potential of *Arthrospira* to adapt to different salinities. Even under a salinity of more than 30 ppt (>0.5 M), *Arthrospira* exhibited a strong growth performance after a time lag of several hours (VONSHAK et al. 1988).

Salinities of more than 8 ppt induced an

increase in filament size. Furthermore, molecular investigations confirmed that the small and large size forms of *Arthrospira fusiformis* from the soda lakes of East Africa are phylogenetically similar (DADHEECH et al. 2010). In this study, we confirmed these findings through molecular phylogenetic analyses of a field sample and the strain isolated from Lake Oloidien (KR2012/1). These results are in agreement with our previous observations on the remarkable conservation in sequences of ITS and *cpcBA*–IGS loci and absence of habitat dependent phylogenetic grouping among *Arthrospira* strains (DADHEECH et al. 2010). Hence filament size has no taxonomic value and currently, the existence of the var. *minor* is doubtful. The populations of small size filaments of *Arthrospira* can be considered as an ecotype growing in alkaline waters of <5 ppt salinity. Furthermore, we did not detect potential cyanotoxin producing genes in the *Arthrospira* strain using different molecular markers. This finding reveals the non-toxic character of the strain (KR2012/1) isolated from Lake Oloidien.

Arthrospira as a food resource for Lesser Flamingos at Lake Oloidien

Does the small size of *Arthrospira* filaments in Lake Oloidien have an impact on food ingestion by Lesser Flamingos? The bill of Lesser Flamingo has fine lamellae, and these lamellae are equipped with rows of fringed platelets which have an average distance of 120 µm. There are about 20 platelets.mm⁻² (JENKIN 1957). The efficiency of this filter apparatus depends on the size of food particles (VARESCHI 1978). Analysis of the stomach contents of Lesser Flamingos revealed that the smallest particles filtered in large quantities were pennate diatoms with a size range of 15–70 µm, while the main food constituent was *Arthrospira* with a size range of 40–800 µm (JENKIN 1957). Hence the small size form of *Arthrospira* from Oloidien fits perfectly into the size range of food particles trapped by the filter apparatus of Lesser Flamingos.

According to BROWN (1959) an adult flamingo with 1.8 kg weight has a minimum daily food requirement of about 180 g FW (= 21.2 g DW). Feeding experiments using caged Lesser Flamingos at Lake Nakuru in the early 1970s revealed that an adult bird must consume much more for its survival (VARESCHI, 1978). A Lesser Flamingo filters 30 l of lake water to ingest 5.6 g DW of *Arthrospira* per hour, which is equal to 72 ± 6.5 g DW per day. The whole flamingo population of about 915,000 birds therefore extracted some 60 t of food algal DW per day, representing 50% – 94% of the daily primary production of the lake or 0.4 – 0.6% of the total algal biomass. We calculated a standing crop of *Arthrospira* in Lake Oloidien to be 236 – 301 t. Using a flamingo density estimate of 200,000 for Lake Oloidien, based on recent estimates, the birds can consume 14.4 t per day. This is 4.8–6.1 % of the standing crop. Hence theoretically, the

flamingos can empty the lake of *Arthrospira* within 16–20 days. However, *Arthrospira* is known for its high growth rate. In culture, both the normal and the small size ecotypes have growth rates of about one doubling per day (GAO & MA 2008). VONSHAK (1997) found doubling times of *Arthrospira* in culture of 11–20 h depending on salinity. However, the calculation of biomass production under field conditions in saline lakes is complicated by the multiple biotic and abiotic interferences (VARESCHI & JACOBS 1985). In natural populations of *Arthrospira* in the soda lake Simbi, an increase of 0.5 mg.l⁻¹ DW per hour based on a standing crop of 100 mg.l⁻¹ DW was measured (MELACK 1979). Applying these findings, rough growth rate estimate for Oloidien of 0.1 mg.l⁻¹ DW h⁻¹ can be calculated based on a standing crop of 20 mg.l⁻¹ DW and a vegetation time of 12 h. This translates in to a daily production of 15.1 t *Arthrospira* in the lake, which is only slightly more than the calculated bird consumption of 14.4 t per day. However, we have to consider that the turnover rate can increase as a result of higher grazing pressure (VARESCHI 1978). In suspension with a lower algal density, the self-shading effects are reduced and this can lead to higher growth rates. Furthermore, the flamingos could utilize additional food resources such as diatoms from the mud and rotifers which occur in high densities in the plankton of Oloidien. Nevertheless, these calculations show that the carrying capacity for flamingos of Lake Oloidien falls in the range between 150,000 and 250,000 birds, hence the observed numbers at the lake did not result in the overexploitation of the food resource. Further investigations are, however, necessary to shed more light on energy flow in the lake including the role of other members of the food chain.

Still, one question on the critical concentration of food algal suspension necessary for normal feeding by flamingos remains unanswered. Based on feeding experiments, VARESCHI (1978, 1982) concluded that flamingos stopped feeding when the biomass of food fell below 100 ± 5 mg.l⁻¹ DW. He therefore concluded that the critical concentration of food algal suspensions necessary for normal feeding by flamingos is 100 ± 5 mg.l⁻¹ DW. We never observed a food algal concentration higher than this critical biomass in Oloidien. The highest value detected during our study was 25 mg.l⁻¹ DW. However, the flamingos did not stop feeding. Comparing data from past studies after converting FW into DW, it is evident that the concentration of *Arthrospira* in the main flamingo lakes during the last decade has been below the critical value of 100 mg.l⁻¹ DW (BALLOT et al. 2004; SCHAGERL & ODUOR 2008; KRIENITZ & KOTUT 2010). It is therefore possible that the flamingos may have changed their feeding behaviour in response to the lower food concentration. An exception to this was Lake Bogoria, which occasionally had a high biomass of *Arthrospira* of about 100 mg.l⁻¹ DW (BALLOT et al.

2004; KAGGWA et al. 2012).

In the light of earlier studies that revealed the presence of cyanotoxins in samples from soda lakes of East Africa (Codd et al. 2003; KRIENITZ et al. 2003; BALLOT et al. 2004, 2005), it is important to note that no cyanotoxins were detected in the phytoplankton of Lake Oloidien, and even the search for toxin–genes in a field clone and two strains isolated turned negative. The emergence of Lake Oloidien as a new feeding habitat for the Lesser Flamingos could not have come at a better time as this took place when the *Arthrospira* populations at two major flamingo–lakes of Elmenteita and Nakuru collapsed in 2010 and 2011 respectively. Lake Oloidien, is a positive development in the history of the Lake Naivasha ecosystem, which has suffered a number of anthropogenic disturbances over the last few decades that include introduction of alien plant and animal species, excessive abstraction of lake water to support geothermal power industry, horticulture and water supply to increasing human settlements in the catchment area (HARPER et al. 2011). The Lesser Flamingos have therefore accepted Lake Oloidien as a new refugium and moved into the lake in high numbers.

ACKNOWLEDGEMENTS

We thank the Government of Kenya for permission to carry out this research (No. MOEST 13/001/31 C 90, and NCST/RRI/12/1/BS/232). We thank the Oloidien Community Boats, members of the initiative “Friends of Flamingos” for assistance in sampling and valuable discussions. PKD was supported by a fellowship of the Leibniz–Institute of Freshwater Ecology and Inland Fisheries. PKD is grateful to Department of Higher Education, Government of Rajasthan, India, for granting permission to carry out research work at IGB, Germany. We thank Monika Degebrot, Monika Papke and Reingard Rossberg for technical support.

REFERENCES

- AL–TEBRINEH, J.; MIHALI, T.K.; POMATI, F. & NEILAN, B.A. (2010): Detection of saxitoxin–producing cyanobacteria and *Anabaena circinalis* in environmental water blooms by quantitative PCR. Appl. – Environ. Microbiol. 76: 7836–7842.
- BALLOT, A.; DADHEECH, P.K. & KRIENITZ, L. (2004): Phylogenetic relationship of *Arthrospira*, *Phormidium* and *Spirulina* strains from Kenyan and Indian waterbodies. – Algological Studies 113: 37–56.
- BALLOT, A.; KOTUT, K.; NOVELO, E. & KRIENITZ, L. (2009): Changes of phytoplankton communities in Lakes Naivasha and Oloidien, examples of degradation and salinization of lakes in the Kenyan Rift Valley. – Hydrobiologia 632: 359–363.
- BALLOT, A.; KRIENITZ, L.; KOTUT, K.; WIEGAND, C.; METCALF, J.S.; CODD, G.A. & PFLUGMACHER, S. (2004): Cyanobacteria and cyanobacterial toxins in three alkaline Rift Valley lakes of Kenya – Lakes Bogoria, Nakuru and Elmenteita. – J. Plankt. Res. 26: 925–35.
- BALLOT, A.; KRIENITZ, L.; KOTUT, K.; WIEGAND, C. & PFLUGMACHER, S. (2005): Cyanobacteria and

- cyanobacterial toxins in the alkaline crater lakes Sonachi and Simbi, Kenya. – *Harmful Algae* 4: 139–150.
- BAURAIN, D.; RENQUIN, L.; GRUBISIC, S.; SCHELDAMAN, P.; BELAY, A. & WILMOTTE, A. (2002): Remarkable conservation of internally transcribed spacer sequences of *Arthrospira* (“*Spirulina*”) (Cyanophyceae, Cyanobacteria) strains from four continents and of recent and 30-year-old dried samples from Africa. – *J. Phycol.* 38: 384–393.
- BBC NATURE (2012): Major flamingo breeding event begins on Lake Natron. <http://www.bbc.co.uk/nature/20271751> (accessed on 26.02.2013).
- BROWN, L. (1959): *The Mystery of the Flamingos*. – Country Life Ltd., London, 116 pp.
- CHILDRESS, B.; HUGHES, B.; HARPER, D.; VAN DEN BOSSCHE, W.V.; BERTHOLD, P. & QUERNER, U. (2006): Satellite tracking documents the East African flyway and key site network of the Lesser Flamingo *Phoenicopterus minor*. – In: BOERE, G.C.; GALBRAITH, C.A. & STROUD, D.A. (eds): *Waterbirds around the world*. – pp. 234–238, The Stationery Office, Edinburgh, UK.
- CHILDRESS, B.; NAGY, S. & HUGHES, B. (Compilers) (2008): International single species action plan for the conservation of the Lesser Flamingo (*Phoeniconaias minor*). – In: CMS Technical Series No. 18. Agreement on the conservation of African–Eurasian Migratory Waterbirds (AEWA) Technical Series No. 34. – 59 pp., Bonn, Germany.
- CODD, G.A.; METCALF, J.S.; MORRISON, L.F.; KRIENITZ, L.; BALLOT, A.; PFLUGMACHER, S.; WIEGAND, C. & KOTUT, K. (2003): Susceptibility of flamingos to cyanobacterial toxins via feeding. – *Vet. Rec.* 152: 722–723.
- DADHEECH, P.K.; BALLOT, A.; CASPER, P.; KOTUT, K.; NOVELO, E.; LEMMA, B.; PRÖSCHOLD, T. & KRIENITZ, L. (2010): Phylogenetic relationship and divergence among planktonic strains of *Arthrospira* (Oscillatoriales, Cyanobacteria) of African, Asian and American origin deduced by 16S–23S ITS and phycocyanin operon sequences. – *Phycologia* 49: 361–372.
- EARTHWATCH INSTITUTE (2006): Lesser Flamingos find new paradise at Kenya’s Oloidien Lake. <http://www.earthwatch.org/europe/newsroom/science/news-3-lesserflamingos.html> (accessed on 31.07.2012)
- EDGAR, R.C. (2004): MUSCLE: multiple sequence alignment with high accuracy and high throughput. – *Nucleic Acids Res.* 32: 1792–1797.
- GAO, K. & MA, Z. (2008): Photosynthesis and growth of *Arthrospira* (*Spirulina*) *platensis* (Cyanophyta) in response to solar radiation, with special reference to its minor variant. – *Environ. & Experiment. Bot.* 63: 123–129.
- HARPER, D.M.; BOAR, R.R.; EVERARD, M. & HICKLEY, P. (eds) (2002): *Lake Naivasha, Kenya*. – Developments in Hydrobiology 168: v–xii, 1–203.
- HARPER, D. M.; CHILDRESS, R. B.; HARPER, M. M.; BOAR, R. R.; HICKLEY, P. H.; MILLS, S. C.; OTIENO, N.; DRANE, T.; VARESCI, E.; NASIRWA, O.; MWATHA, W. E.; DARLINGTON, J. P. E. C. & ESCUTÉ-GASULLA, X. (2003): Aquatic biodiversity and saline lakes: Lake Bogoria National Reserve, Kenya. – *Hydrobiologia* 500: 259–276.
- HARPER, D.M.; MORRISON, E.H.J.; MACHARIA, M.M.; MAVUTI, K.M. & UPTON, C. (2011): *Lake Naivasha, Kenya: ecology, society and future*. – *Freshw. Rev.* 4: 89–114.
- HARPER, D.M., MUCHANE, M., KIMANI, D.K. & MWINAMI, T. (2006): Thousands of Lesser Flamingos at Lake Naivasha? – *Scopus* 26, 8–10.
- HEPPERLE, D. (2008): Align, Multisequence Alignment Editor ver. 05/2008 SequentiX – Digital DNA Processing, Klein Raden, Germany, www.sequentix.de.
- JENKIN, P.M. (1957): The filter feeding and food of flamingos. (Phoenicopter). – *Philos. Trans. R. Soc. B* 240: 401–493.
- JUNGBLUT, A.D. & NEILAN, B.A. (2006): Molecular identification and evolution of the cyclic peptide hepatotoxins, microcystin and nodularin, synthetase genes in three orders of cyanobacteria. – *Arch. Microbiol.* 185: 107–114.
- KAGGWA, M.N.; GRUBER, M.; ODUOR, S.O. & SCHAGERL, M. (2012): A detailed time series assessment of the diet of Lesser Flamingos: further explanation for their itinerant behaviour. – *Hydrobiologia* 710: 83–93.
- KALFF, J. & WATSON, S. (1986): Phytoplankton and its dynamics in two tropical lakes: a tropical and temperate zone comparison. – *Hydrobiologia* 138: 161–176.
- KRIENITZ, L. & KOTUT, K. (2010): Fluctuating algal food populations and the occurrence of Lesser Flamingos (*Phoeniconaias minor*) in three Kenyan Rift Valley lakes. – *J. Phycol.* 46: 1088–1096.
- KRIENITZ, L. & WIRTH, M. (2006): The high content of polyunsaturated fatty acids in *Nannochloropsis limnetica* (Eustigmatophyceae) and its implication for food web interactions, freshwater aquaculture and biotechnology. – *Limnologia* 36: 204–210.
- KRIENITZ, L.; BALLOT, A.; KOTUT, K.; WIEGAND, C.; PÜTZ, S.; METCALF, J. S.; CODD, G.A. & PFLUGMACHER, S. (2003): Contribution of hot spring cyanobacteria to the mysterious deaths of Lesser Flamingos at Lake Bogoria, Kenya. – *FEMS Microbiol. Ecol.* 1437: 141–148.
- KRIENITZ, L.; DADHEECH, P.K. & KOTUT, K. (2013): Mass developments of the cyanobacteria *Anabaenopsis* and *Cyanospira* (Nostocales) in the soda lakes of Kenya: ecological and systematic implications. – *Hydrobiologia* 703: 79–93.
- MAINA, C. & KYUNGU, S. (eds.) (2009): *The African Waterfowl Census*. – Museum News, March–June 2009, 8.
- MELACK, J.M. (1979): Photosynthesis and growth of *Spirulina platensis* (Cyanophyta) in an equatorial lake (Lake Simbi, Kenya). – *Limnol. Oceanogr.* 24: 753–760.
- NASIRWA, O.; OWINO, A.; MUCHAI, M. & NDANG’ANG’A, K. (2007): Assessing trends in waterbird numbers in major Kenyan wetland sites (1991–2004). – *Ostrich* 78: 551–561.
- NEILAN, B.A.; DITTMANN, E.; ROUHIAINEN, L.; BASS, A.; SCHAUB, V.; SIVONEN, K. & BÖRNER, T. (1999): Nonribosomal peptide synthesis and toxigenicity of cyanobacteria. – *J. Bacteriol.* 181: 4089–4097.
- OPTICOUNT (2008): – http://science.do-mix.de/software_opticount.php
- OWINO, A.; OYUGI, J.O.; NASIRWA, O.O. & BENNUN, L.A. (2001): Patterns of variation in waterbird numbers on four Rift Valley lakes in Kenya. – *Hydrobiologia* 458: 45–53.
- RANTALA-YLINEN, A.; KANA, S.; WANG, H.; ROUHIAINEN, L.;

- WAHLSTEN, M.; RIZZI, E.; BERG, K.; GUGGER, M. & SIVONEN, K. (2011): Anatoxin-a synthetase gene cluster of the cyanobacterium *Anabaena* sp. strain 37 and molecular methods to detect potential producers. – *Appl. Environ. Microb.* 77: 7271–7278.
- RICH, F. (1932): Reports on the Percy Sladen Expedition to some Rift Valley Lakes in Kenya in 1929. – IV. Phytoplankton from the Rift Valley Lakes in Kenya. 233–262. – *Ann. Magaz. Nat. Hist.*, ser. 10, 10: 233–262.
- SCHAGERL, M. & ODUOR, S.O. (2008): Phytoplankton community relationship to environmental variables in three Kenyan Rift Valley saline-alkaline lakes. – *Marin. Freshw. Res.* 59:125–36.
- STANDARD MEDIA (2012): Some of the 1.5 million flamingoes fly around Lake Oloidien. <http://www.standardmedia.co.ke/?id=2000027095&cid=468&story>
- TAMURA, K.; PETERSON, D.; PETERSON, N.; STECHER, G.; NEI, M. & KUMAR, S. (2011): MEGA5: Molecular evolutionary genetics Analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. – *Mol. Biol. Evol.* 28: 2731–2739.
- TUITE, C. H. (1981): Standing crop densities and distribution of *Spirulina* and benthic diatoms in East African alkaline saline lakes. – *Freshw. Biol.* 11: 345–360.
- TUITE, C. H. (2000): The distribution and density of Lesser Flamingos in East Africa in relation to food availability and productivity. – *Waterbirds (Spec. Publ.)* 23: 52–63.
- UTERMÖHL, H. (1958): Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. – *Mitt. Int. Verein. Limnol.* 9: 1–38.
- VARESCHI, E. (1978): The ecology of Lake Nakuru (Kenya) I. Abundance and feeding of the Lesser Flamingo. – *Oecologia* 32:11–35.
- VARESCHI, E. (1982): The ecology of Lake Nakuru (Kenya) III. Abiotic factors and primary production. – *Oecologia* 55:81–101.
- VARESCHI, E. & JACOBS, J. (1985): The ecology of Lake Nakuru VI. Synopsis of production and energy flow. – *Oecologia* 65:412–24.
- VERSCHUREN, D.; COCQUYT, C.; TIBBY, J.; ROBERTS, C.N. & LEAVITT, P.R. (1999): Long-term dynamics of algal and invertebrate communities in a small, fluctuating tropical soda lake. – *Limnol. Oceanogr.* 44: 1216–1231.
- VERSCHUREN, D.; TIBBY, J.; SABBE, K. & ROBERTS, N. (2000): Effects of depth, salinity, and substrate on the invertebrate community of a fluctuating tropical lake. – *Ecology* 81: 164–182.
- VONSHAK, A. (1997): *Spirulina*: Growth, physiology and biochemistry. – In: VONSHAK, A. (ed.): *Spirulina platensis (Arthrospira) physiology, cell biology and biotechnology*. – pp. 43–65, Taylor & Francis, London, UK.
- VONSHAK, A. & RICHMOND, A. (1988): Mass production of the blue-green algae *Spirulina*, an overview. – *Biomass* 15: 233–247.
- VONSHAK, A.; GUY, R. & GUY, M. (1988): The response of the filamentous cyanobacterium *Spirulina platensis* to salt stress. – *Arch. Microbiol.* 150: 417–420.